

REEF EXPERT EVALUATION INVITATION

Dual-Layer Stratified Nanobubble Shading and Oxygen Support for Coral Reef Thermal Stress Mitigation

Origin: [ReefHealer](#) service system by [Alarivean](#)

Status: Core systems at Technology Readiness Level 8 (TRL8)

Date: October 2026

Target Community: Reef Stewardship Experts

Location Context: Reef Lagoons / Nurseries

Framework: Aquatic Prosperity as a Service (APaaS)

1. Executive Summary & Context

As marine heatwaves intensify globally, traditional macroscopic engineering interventions prove insufficient or too ecologically dangerous for fragile reef protection. Alarivean, whose core systems operate at Technology Readiness Level 8 (TRL8), introduces a highly localized, mechanically generated, pulse-managed intervention framework that adds no surfactant, designed specifically to protect delicate habitats during acute summer thermal stress events without global environmental side effects. Any carbonate reagent is evaluated as a separate, measured trial arm. TRL8 covers our core systems: the proven vessel delivery capability and the mechanisms with a solid scientific basis. The less certain and research-stage items, such as antioxidant support, sit within our service model, which is contracted to outcomes rather than to static equipment or outputs. That lets us keep innovating, and we will always do so under independent scientific leadership and accountability.

Moving away from speculative atmospheric modifications, our operational architecture centers on the **ReefHealer** platform deployed via quiet, gentle, shallow-draft electric **SeaBreather** vessel variants. These systems process local waters through an internal loop, applying localized dual-layer nanobubble infusion before releasing them cleanly back into target columns. No chemical stabilizers are added. How long the bubble field persists in site seawater, how it dissipates once deployment concludes, and any physical or chemical footprint on the ecosystem are measured rather than assumed.

To support early academic validation and minimize operational overhead, Alarivean introduces a **Minimum Viable System (MVS)** configuration for field calibration trials, which begin with a measurement-only baseline on treated and control plots before any dosing: residence time and site currents, pre-dawn dissolved oxygen (DO), total dissolved gas (TDG), the carbonate system, hydrogen peroxide (H₂O₂), and PAR and UV at coral depth. Oxygen and carbonate are then calibrated in a poorly flushed lagoon cell, and shading is compared with shade cloth in a nursery. This localized baseline allows stewardship teams to monitor and verify fine-scale column interactions, bubble kinetics, and target biological indicators before scaling up to full logistics networks.

2. The Hybrid Architectural Hypothesis

We invite reef stewardship experts to collaborative evaluation of a multi-tiered mitigation architecture that decouples stress remediation into two independent vectors (shading and oxygen), plus an optional carbonate arm trialed separately, with shading applied during peak daylight hours and oxygen dosed at night as well as by day:

- **The Surface Shading Layer (Natural Air):** Injection of natural air nanobubbles into the upper surface column during peak daylight hours. The surface bubble field shades the water beneath it by scattering incoming solar and ultraviolet (UV) radiation. We are compiling measurements of how much shade it provides in which contexts, as PAR and UV at coral depth alongside coral responses, with light starvation tracked at the same time. Bubble size distribution, number density and lifetime are measured in site seawater.

- **The Dissolved Oxygen Support Layer (Sub-Surface Benthic Loop):** Infusion of pure oxygen (O₂) nanobubbles into the water column above the reef architecture, intended to counter hypoxia during heat stress driven by thermal stratification and high biological oxygen demand (BOD). Reef hypoxia is mainly nocturnal and worst in stagnant lagoon water, so we dose at night as well as by day, under a total dissolved gas (TDG) cap.
- **Optional Carbonate Arm (Sodium Carbonate):** Sodium carbonate (soda ash, Na₂CO₃) delivered pre-dissolved, at a discharge depth set for the target and a dose rate stated per site and logged per lot, capped at +0.1 pH at the target, with a discharge-point pH ceiling; the plume is diluted below an aragonite saturation (Ω_{arag}) of about 5 within a stated mixing distance, a threshold from lime-addition experiments to be confirmed for sodium carbonate (Moras et al., 2022), and sodium carbonate is not dosed where ambient water already exceeds it. It offers carbonate chemistry that may support calcification where measured. It is added as a solution, not as powder or grains, and is not used for shading; the plume is checked for calcium carbonate precipitation, and dosing stops if it is detected. Our coral mode doses no peroxide or other oxidant, and residual oxidant is measured independently.

Core Thesis: By decoupling the shading mechanism (natural air) from the respiration support mechanism (pure O₂), and timing each to its hours of peak stress (midday light for shading, the nighttime oxygen minimum for respiration support), we aim to lower light stress during heat events while sustaining reef respiration, and we track the trade-offs of prolonged light starvation and hypoxia.

3. Target Themes & Key Questions

Alarivean invites reef stewardship experts to co-design and independently audit verification tracks exploring several core themes:

Theme 1: Structural Effects of Elevated Dissolved Oxygen (DO)

- How do localized pure O₂ nanobubbles affect the oxidative stress thresholds of coral holobionts? Does localized O₂ enrichment raise or lower holobiont reactive oxygen species (ROS) under heat and high light, does the answer change once symbiont photosynthesis fails, and which moves the pre-dawn oxygen minimum more: daytime or nighttime dosing?
- What are the micro-respiratory dynamics of cryptic reef invertebrates, bioeroders, and endolithic communities when subjected to localized hyperoxia during extreme thermal anomalies?

Theme 2: Intermittent Shading & Carbonate Chemistry

- What are the precise light starvation thresholds of diverse reef communities when subjected to temporary shading by a surface air-bubble field of measured size, density and lifetime, and how does that shade, measured as PAR and UV at coral depth, compare with shade cloth in a nursery? Separately, does a sodium carbonate dose change calcification at reef-plausible doses, and does raised oxygen cancel that gain (Wijgerde et al., 2014)?
- How do these thresholds shift across different operational scenarios, including variations in diurnal loops, seasonal extremes, depth profiles, and distinct taxonomic groups (e.g., branching vs. massive scleractinians, octocorals)?
- How much treated water, and what dose, reaches targets the vessels do not pass over, such as patch-reef tops, shallow nursery margins and reef flats, and which tracer best establishes the dose at the target?

Theme 3: Algal & Nutrient Loads in HAB and Water-Treatment Mode, Run Clear of Live Coral

- By utilizing the vessel-internal loop and pass-through Advanced Oxidation Processes (AOP) within onboard *HABslayer* and *ChemSlayer* modules, what are the precise target extraction ratios required to strip excessive nutrient burdens and suppress opportunistic macroalgal overgrowth?
- How can we tune these internal AOPs to act not like an indiscriminate chemical hand sanitizer that leaves a biological vacuum, but like a gentle, targeted preventative wash that knocks down harmful imbalances while letting remaining native biota infill healthy immune defenses and restore natural dominance?
- What residual-oxidant limits at the mixing-zone edge, stand-off distances, current-direction rules and quench chemistry keep HAB-mode discharge from reaching live coral, including during spawning, and what brominated byproducts form in seawater under the AOP chemistry we use?

Theme 4: Antioxidant Support (Research Only)

- Is there a safe and effective antioxidant regime for corals in open water? Ascorbate (vitamin C) generates hydrogen peroxide in seawater and consumes alkalinity, and mortality was reported at laboratory doses in an anemone study (Dungan et al., 2022a). Mannitol, probiotics and thermal preconditioning are alternative routes (Dungan et al., 2022a, 2022b; Majerová & Drury, 2022). This is a research question; we propose no dose.

4. Bounded Trial Protocols & Live Safeguards

To ensure uncompromised scientific integrity and complete alignment with local environmental authorities, all Alarivean APaaS (Aquatic Prosperity as a Service) calibration trials operate under strict governance parameters:

- **Dual-Key Data Feeds:** Independent overseeing academic institutions hold a live, unedited stream of all raw sensor telemetry and water quality datasets in real time.
- **Absolute Infraction Halts:** Collaborating principal scientists maintain absolute, unilateral veto authority to immediately shut down or pause all vessel system operations upon any suspected boundary anomaly.
- **Surfactant Prohibition:** All trials utilize strictly mechanical, surfactant-free generation methods: no surfactants are added. Bubble size distribution and lifetime in site seawater are measured, not assumed; natural surface-active material in reef water can itself stabilize bubbles (Johnson & Cooke, 1981), so 'surfactant-free' describes what we add, not the bubbles.
- **Oxidant Separation:** Coral mode runs at a zero oxidant set-point: no peroxide or other oxidant is dosed. AOP modules run only in HAB or water-treatment mode, at a stated stand-off from live coral, down-current of it and outside spawning windows, within residual-oxidant limits at the mixing-zone edge, with a named quench, an independent assay and dose logs shared with the overseeing institutions.
- **Reagents, Permits & Gas Load:** A certificate of analysis for each sodium carbonate lot; the discharge permits named for each site; and a total dissolved gas cap for pure-O₂ injection.
- **Navigation Safeguards:** SeaBreather vessels operate strictly using shallow-draft electric propulsion and will never navigate directly over or make physical contact with shallow reef structures; all processing occurs via pass-through flow adjacent to or above safe bathymetric zones. Vessels operate only where the water is at least 2 to 3 m deep, so the hull never comes close enough to live coral to risk contact. Discharge depth is set by the service outcome, anywhere from just below the surface downward. Targets the vessels do not pass over, such as patch-reef tops, shallow nursery margins and reef flats, are reached by treated water rather than treated from above: a dye or other tracer establishes the dose that arrives, with monitoring stations at the target and down-current. Treated water that drains over a reef flat at low tide is monitored there for dissolved

oxygen, TDG, pH and Ω arag against the flat's own natural range, with a stop rule, for example TDG or Ω arag outside the flat's baseline range.

Interested verification teams, scientific labs, and field stations are invited to submit situation reports or joint protocol designs directly to Alarivean (Alarivean.com).

5. Draft Framework

This section reviews the current state of coastal marine science regarding targeted environmental modifications inside delicate coral reef lagoons. Rather than addressing broad planetary geoengineering, it synthesizes the known biological and physical impacts of localized, high-density interventions designed to alleviate thermal and radiative anomalies using Alarivean's ReefHealer and SeaBreather platforms. Anthropogenic warming has drastically elevated the frequency and intensity of marine heatwaves, pushing coral holobionts past their evolutionary tipping points (Hughes et al., 2017; Hoegh-Guldberg et al., 2007; Altieri et al., 2017). Traditional marine protected areas provide critical spatial defensive framing but remain fundamentally passive during acute atmospheric anomalies (Altieri et al., 2017; Nelson & Altieri, 2019). The Alarivean architecture establishes a pulse-managed, highly targeted intervention framework deployed via quiet, gentle, shallow-draft electric SeaBreather vessel variants. These platforms utilize vessel-internal loops to condition lagoon water via localized nanobubble infusion, avoiding direct structural reef contact while optimizing the microenvironment scale.

Intervention Vector	Verified Scientific Certainty (Solid)	Verification Opportunities (Less Certain)
1. Dissolved Oxygen Elevation	Thermal stress induces water column stratification and acute nocturnal hypoxia, accelerating coral tissue necrosis (Altieri et al., 2017; Nelson & Altieri, 2019). Localized tissue-surface anoxia preceded rapid coral tissue disintegration under acute heat stress (Dhillon et al., 2026). In one laboratory study, bulk oxygen supersaturation (hyperoxia at 10 mg O ₂ L ⁻¹) raised the median lethal thermal threshold (LT ₅₀) by 0.2°C in one of three reef-building species (<i>Galaxea fascicularis</i>) under a single 10-hour ramp to 40°C (Parry et al., 2025). Deoxygenation lowers the bleaching threshold (Alderdice et al., 2022), and reef hypoxia is mainly nocturnal (Pezner et al., 2023).	Physiological limits of induced hyperoxia in cryptic reef organisms and micro-scale endolithic communities; mapping the optimal balance of boundary-layer artificial ventilation across distinct hydrodynamic fields to alleviate severe respiratory supply-demand mismatches.
2. Surface Shading & Carbonate Chemistry	Shading reduced heat-driven bleaching in aquarium and nursery trials (Butcherine et al., 2023; Ruiz-Diaz et al., 2025). Alkalinity addition raised net calcification on a reef flat (Albright et al., 2016) and calcification in <i>Acropora cervicornis</i> (Cooke et al., 2026). How much shade the surface bubble field delivers in each context: we are compiling PAR and UV at coral depth against bubble density, lifetime and water clarity, with coral responses. Whether carbonate gains hold under heat stress and alongside raised oxygen (Wijgerde et al., 2014).	Ecosystem light starvation risks: defining the exact balance where shading benefits are outweighed by light reduction across diurnal loops, seasonal extremes, and distinct depth or taxonomic profiles (branching acroporids vs. massive scleractinians vs. octocorals).
3. Nutrient & Algal Reduction (HAB Mode)	High organic and nutrient loads fuel opportunistic macroalgal blooms that smother corals and deplete oxygen layers (Seitzinger et al., 2010; Fabricius, 2005). Vessel-internal pass-through loop processing running Advanced Oxidation Processes (AOP) avoids physical	Eco-tuning thresholds: identifying the exact drawdown ratios required to allow native ecosystems to self-stabilize and reclaim dominance from turf algae without

Intervention Vector	Verified Scientific Certainty (Solid)	Verification Opportunities (Less Certain)
	mass displacement. AOP runs only in HAB or water-treatment mode, clear of live coral.	over-sterilizing the column or stripping beneficial native microbiota and larval pathways.

6. Detailed Synthesis of Intervention Vectors

Vector 1: Column Dissolved Oxygen Dynamics

Thermal stress does not operate in an environmental vacuum. As sea surface temperatures climb, oxygen solubility in the upper water column falls, while biological oxygen demand (BOD) simultaneously accelerates (Vaquer-Sunyer & Duarte, 2011; Altieri et al., 2017; Parry et al., 2025). In shallow, poorly flushed coral lagoons, this physics-driven reduction in solubility is severely compounded by vertical thermal stratification and heavy nocturnal community respiration, generating acute benthic hypoxia (Altieri et al., 2017; Adelson et al., 2022; Nelson & Altieri, 2019). This hypoxia is mainly nocturnal (Pezner et al., 2023), so we dose oxygen at night as well as by day. In laboratory microsensor work, oxygen depletion at the tissue–water interface preceded the onset of rapid coral tissue disintegration—a catastrophic structural collapse distinct from classical bleaching (Dhillon et al., 2026). Short-term anoxia alone is sufficient to trigger this tissue disintegration phenotype, consistent with mass-transfer limitation across a stagnant concentration boundary layer (CBL) creating a localized metabolic crisis during heatwaves (Dhillon et al., 2026). High flow rates thin the CBL, acting as a powerful hydrodynamic countermeasure (Dhillon et al., 2026). In one laboratory study, bulk seawater oxygen supersaturation (hyperoxia at 10 mg O₂ L⁻¹) raised the median lethal thermal threshold (LT₅₀) by 0.2°C in one of three reef-building species (*Galaxea fascicularis*, a single 10-hour ramp), consistent with a passive diffusion buffer once microalgal photosynthetic oxygen production collapses (Parry et al., 2025). In laboratory studies without heat stress, raised oxygen reduced calcification gains and the photosynthetic benefit of flow (Wijgerde et al., 2014; Mass et al., 2010), so oxygen is dosed to a stated cap. Crucially, oxygen availability also modulates the metabolic capacity of associated reef fish, such as how oxygen availability modulates clownfish larvae growth under thermal stress (Arossa et al., 2025). The ReefHealer platform is designed around these dynamics, infusing pure O₂ nanobubbles into a benthic loop to counter local hypoxia, especially at night and in stagnant lagoon water.

Vector 2: Localized Surface Shading

Severe mass bleaching events are rarely driven by elevated temperature alone; the cumulative destructive impact is multiplied by high downwelling solar irradiance, specifically Photosynthetically Active Radiation (PAR) and shortwave ultraviolet (UV) spectra (Hoegh-Guldberg et al., 2007; Hughes et al., 2017). Excessive photon fluxes overload the light-harvesting complex of symbiotic dinoflagellates (Symbiodiniaceae), damaging their photosynthetic machinery and generating high concentrations of toxic reactive oxygen species (ROS) that prompt host cells to sever the symbiosis (Iglesias-Prieto et al., 1992; Warner et al., 1999; Hoegh-Guldberg et al., 2007). A near-surface field of natural air bubbles shades the water beneath it: the bubbles scatter incoming light diffusely, so less PAR and UV reaches the reef. We are compiling measurements of how much shade the field provides in which contexts (water clarity, depth, bubble density and generator throughput), alongside coral responses, with light starvation tracked at the same time and shade cloth as the comparison in nurseries (Butcherine et al., 2023; Ruiz-Diaz et al., 2025). Carbonate chemistry is a separate, optional arm. Sodium carbonate is added as a solution, not as powder or grains, and is not used for shading; it offers carbonate chemistry that may support calcification where measured (Albright et al., 2016; Cooke et al., 2026). The dose is capped at +0.1 pH at the target, with a discharge-point

pH ceiling; the plume is diluted below an aragonite saturation (Ω_{arag}) of about 5 within a stated mixing distance, a threshold from lime-addition experiments to be confirmed for sodium carbonate (Moras et al., 2022), and we do not dose where ambient water already exceeds it. Visible whitening near the injection point would indicate calcium carbonate precipitation, which consumes the added alkalinity and reduces light at coral depth; we monitor for it, and dosing stops if it is detected (Moras et al., 2022; Hartmann et al., 2023). Our coral mode doses no peroxide or other oxidant. The data we are compiling cover bubble size distribution and lifetime in site seawater, the reduction in PAR and UV at coral depth with the coral response, and light starvation risks across diurnal loops, seasonal extremes, and distinct taxonomic depth profiles (Hoegh-Guldberg et al., 2007).

Vector 3: Nutrient Extraction and Algal Suppression (HAB Mode)

Degraded coastal reef environments are frequently driven into toxic ecological phase shifts, where land-based nutrient enrichment and organic loading tip the competitive balance away from calcifying scleractinian corals and toward fast-growing turf and macroalgal communities (Seitzinger et al., 2010; Fabricius, 2005; Burkepile & Hay, 2006). These opportunistic algal mats form dense physical barriers that prevent coral larval settlement, trap sediment, block localized light transmission, and accumulate organic materials that exacerbate localized boundary-layer oxygen depletion (Fabricius, 2005; Burkepile & Hay, 2006; Hughes et al., 2010). The SeaBreather platform addresses these organic and chemical crises by utilizing gentle, vessel-internal containment loops running advanced onboard oxidation processes (AOP), featuring specialized HABslayer and ChemSlayer operational modules, which run only in HAB and water-treatment mode, clear of live coral, under the oxidant separation conditions in our trial safeguards. This pass-through architecture isolates and treats water masses internally, ensuring that no physical mass displacement occurs within the habitat footprint. The core verification gap within this vector focuses on defining precise eco-tuning thresholds. The internal AOP reactor must not operate like an indiscriminate chemical hand sanitizer that eliminates all organic mass, which would generate a biological vacuum and strip out beneficial pelagic microbiota, planktonic larvae, and necessary low-level nutrients (Seitzinger et al., 2010; Fabricius, 2005; Hughes et al., 2010). Instead, it functions as a gentle, targeted preventative wash. The fundamental scientific objective is to determine the exact extraction ratios required to drop opportunistic turf algal growth rates just enough to allow native communities to self-stabilize, infill critical ecological niches, and naturally restore long-term benthic dominance.

7. Operational In-Situ Monitoring Matrix

To ensure absolute safety, multi-lateral scientific transparency, and clear isolation of variables during a localized calibration trial using an Alarivean Minimum Viable System (MVS), field monitoring must track key diagnostic metrics across three distinct sequential phases (Altieri et al., 2017; Hoegh-Guldberg et al., 2007; Dhillon et al., 2026). Continuous microsensor and optical arrays will map boundary layer microenvironments to validate system performance in real time. The pre-deployment baseline is measurement-only, on treated and control plots, and no dosing begins until it is complete.

Operational Phase	Core Target Parameter	Measurement Methodology / Instrumentation
Pre-Deployment Baseline	Baseline PAR & Vertical Attenuation	Deployment of multi-spectral radiometers across reef bathymetry zones to map ambient light transmission curves, including UV-A and UV-B, on treated and control plots.

Operational Phase	Core Target Parameter	Measurement Methodology / Instrumentation
Pre-Deployment Baseline	Ambient Dissolved Oxygen & Nutrients	High-frequency electrochemical loggers & optical DO microsensors paired with colorimetric assays for baseline N/P profiles (Altieri et al., 2017; Dhillon et al., 2026).
Pre-Deployment Baseline	Pre-Dawn DO, TDG & Carbonate System	Measurement-only: pre-dawn DO minimum, and DO and temperature at colony height; total dissolved gas; total alkalinity, pH and Ω_{arag} ; H ₂ O ₂ by an H ₂ O ₂ -specific method.
Pre-Deployment Baseline	Residence Time, Currents & Water Clarity	Current meters and drifters for residence time and site currents; K _d (PAR), particulate backscatter, black-disk visibility and bottom type at each target.
Active Treatment Phase	Temperature at Coral Depth	Benthic thermistor arrays logging temperature at coral depth and colony height on treated and control plots, reported as a measurement.
Active Treatment Phase	Photosynthetic Efficiency (F _v /F _m)	In-situ Pulse-Amplitude Modulation (PAM) fluorometry on live coral colonies to evaluate operational quantum yield of Photosystem II (Hoegh-Guldberg et al., 2007).
Active Treatment Phase	PAR & UV at Coral Depth	Radiometers logging PAR, UV-A and UV-B at coral depth as daily light integral and peak irradiance on treated and control plots, with shade cloth as the nursery comparator; black-disk visibility and K _d (PAR) are logged during treatment.
Active Treatment Phase	Dose at the Target	Dye or another conservative tracer released with treatment, with stations at the target and down-current, to establish the dose reaching targets the vessels do not pass over.
Active Treatment Phase	Carbonate System & Precipitation	Total alkalinity, dissolved inorganic carbon, pH and Ω_{arag} at the discharge point, at the target and in O ₂ -only plots; turbidity, filtered samples and particulate inorganic carbon to detect calcium carbonate precipitation (Schulz et al., 2023); sediment traps at the target.
Active Treatment Phase	Gas Load & Oxidant Residuals	TDG at depth and O ₂ saturation at the coral surface; a coral ROS or oxidative-damage endpoint; in HAB mode only, H ₂ O ₂ , total residual oxidant by DPD and brominated byproducts by GC, with method and detection limit named.
Active Treatment Phase	Microbial Equilibrium Retention	High-throughput flow cytometry & eDNA pass-through loop sampling to monitor native microbiota stability and planktonic larval pathways.
Post-Recovery Phase	Long-term Tissue & Calcification Health	Skeletal growth micro-CT scanning of structural core samples paired with photographic transect monitoring over a 30–90 day tracking loop.

8. Peer-Reviewed Academic References

- Adelson, A. E., Altieri, A. H., Boza, X., et al. (2022). Seasonal hypoxia and temperature inversions in a tropical bay. *Limnology and Oceanography*, 67(10), 2174–2189. <https://doi.org/10.1002/lno.12196>
- Albright, R., et al. (2016). Reversal of ocean acidification enhances net coral reef calcification. *Nature*, 531(7594), 362–365. <https://doi.org/10.1038/nature17155>
- Alderdice, R., et al. (2022). Deoxygenation lowers the thermal threshold of coral bleaching. *Scientific Reports*, 12, 18273. <https://doi.org/10.1038/s41598-022-22604-3>
- Altieri, A. H., et al. (2017). Tropical dead zones and mass mortalities on coral reefs. *Proceedings of the National Academy of Sciences*, 114(14), 3660–3665.
- Arossa, S., Steckbauer, A., Klein, S. G., Parry, A. J., Alva Garcia, J. V., Alamoudi, T., Pluma, N., Frappi, S., & Duarte, C. M. (2025). Oxygen availability modulates clownfish larvae responses to acute thermal stress in experimental settings. *Proceedings of the Royal Society B*, 292(2058), 20252201. <https://doi.org/10.1098/rspb.2025.2201>

- Burkepile, D. E., & Hay, M. E. (2006). Herbivore vs. nutrient control of marine primary producers: context-dependent effects. *Ecology*, 87(12), 3128–3139.
- Butcherine, P., et al. (2023). Intermittent shading can moderate coral bleaching on shallow reefs. *Frontiers in Marine Science*, 10, 1162896. <https://doi.org/10.3389/fmars.2023.1162896>
- Cooke, K. M., et al. (2026). Alkalinity enrichment stimulates calcification and linear extension in *Acropora cervicornis*. *Scientific Reports*, 16, 14512. <https://doi.org/10.1038/s41598-026-44817-6>
- Dhillon, M. S., Klein, S. G., Maity, D. K., Parry, A. J., Moret, A., Kushwaha, A. K., Dighe, S., Pachherres, C. O., Köhl, M., Truscott, T., Dellisanti, W., Aranda, M., & Duarte, C. M. (2026). Oxygen Deprivation Implicated in Rapid Coral Mortality Under Acute Heating Events. *Global Change Biology*, 32(8), e71030. <https://doi.org/10.1111/gcb.71030>
- Dungan, A. M., et al. (2022a). Lack of evidence for the oxidative stress theory of bleaching in the sea anemone, *Exaiptasia diaphana*, under elevated temperature. *Coral Reefs*, 41(4), 1161–1172. <https://doi.org/10.1007/s00338-022-02251-w>
- Dungan, A. M., Hartman, L. M., Blackall, L. L., & van Oppen, M. J. H. (2022b). Exploring microbiome engineering as a strategy for improved thermal tolerance in *Exaiptasia diaphana*. *Journal of Applied Microbiology*, 132(4), 2940–2956. <https://doi.org/10.1111/jam.15465>
- Fabricius, K. E. (2005). Effects of terrestrial runoff on the ecology of corals and coral reefs: review and synthesis. *Marine Pollution Bulletin*, 50(2), 125–146.
- Hartmann, J., et al. (2023). Stability of alkalinity in ocean alkalinity enhancement (OAE) approaches – consequences for durability of CO₂ storage. *Biogeosciences*, 20(4), 781–802. <https://doi.org/10.5194/bg-20-781-2023>
- Hoegh-Guldberg, O., et al. (2007). Coral reefs under rapid climate change and ocean acidification. *Science*, 318(5857), 1737–1742.
- Hughes, T. P., et al. (2017). Global warming and recurrent mass bleaching of corals. *Nature*, 543(7645), 373–377.
- Hughes, T. P., et al. (2010). Rising to the challenge of sustaining coral reef resilience. *Trends in Ecology & Evolution*, 25(11), 633–642.
- Iglesias-Prieto, R., et al. (1992). Photosynthetic response to elevated temperature in the symbiotic dinoflagellate *Symbiodinium microadriaticum*. *Proceedings of the National Academy of Sciences*, 89(21), 10302–10305.
- Johnson, B. D., & Cooke, R. C. (1981). Generation of stabilized microbubbles in seawater. *Science*, 213(4504), 209–211. <https://doi.org/10.1126/science.213.4504.209>
- Majerová, E., & Drury, C. (2022). Thermal preconditioning in a reef-building coral alleviates oxidative damage through a BI-1-mediated antioxidant response. *Frontiers in Marine Science*, 9, 971332. <https://doi.org/10.3389/fmars.2022.971332>
- Mass, T., et al. (2010). Flow enhances photosynthesis in marine benthic autotrophs by increasing the efflux of oxygen from the organism to the water. *Proceedings of the National Academy of Sciences*, 107(6), 2527–2531. <https://doi.org/10.1073/pnas.0912348107>
- Moras, C. A., Bach, L. T., Cyronak, T., Joannes-Boyau, R., & Schulz, K. G. (2022). Ocean alkalinity enhancement – avoiding runaway CaCO₃ precipitation during quick and hydrated lime dissolution. *Biogeosciences*, 19(15), 3537–3557. <https://doi.org/10.5194/bg-19-3537-2022>
- Nelson, H. R., & Altieri, A. H. (2019). Oxygen: the universal currency on coral reefs. *Coral Reefs*, 38(2), 177–198. <https://doi.org/10.1007/s00338-019-01765-0>
- Parry, A. J., Klein, S. G., Steckbauer, A., Dhillon, M. S., Muniz Barreto, M., Golding, V. C., Re, E., Arossa, S., Aranda, M., & Duarte, C. M. (2025). Oxygen supersaturation could protect reef-building corals against acute thermal stress. *Global Change Biology*, 31, e70586. <https://doi.org/10.1111/gcb.70586>
- Pezner, A. K., et al. (2023). Increasing hypoxia on global coral reefs under ocean warming. *Nature Climate Change*, 13(4), 403–409. <https://doi.org/10.1038/s41558-023-01619-2>
- Ruiz-Diaz, C. P., et al. (2025). Advancing coral conservation: shading strategies to mitigate heat and light stress in *Acropora cervicornis* amid intensifying climate challenges. *Conservation Evidence Journal*, 22, 45–56. <https://doi.org/10.52201/CEJ22/QWAT5710>
- Schulz, K. G., Bach, L. T., & Dickson, A. G. (2023). Seawater carbonate chemistry considerations for ocean alkalinity enhancement research: theory, measurements, and calculations. *State of the Planet*, 2-oae2023, 2. <https://doi.org/10.5194/sp-2-oae2023-2-2023>
- Seitzinger, S. P., et al. (2010). Global river nutrient export: A scenario analysis of past and future trends. *Global Biogeochemical Cycles*, 24(4), GBOA08.

- Vaquer-Sunyer, R., & Duarte, C. M. (2011). Temperature effects on oxygen thresholds for hypoxia in marine benthic organisms. *Global Change Biology*, 17(5), 1788–1797. <https://doi.org/10.1111/j.1365-2486.2010.02343.x>
- Warner, M. E., Fitt, W. K., & Schmidt, G. W. (1999). Damage to photosystem II in symbiotic dinoflagellates: a determinant of coral bleaching. *Proceedings of the National Academy of Sciences*, 96(14), 8007–8012. <https://doi.org/10.1073/pnas.96.14.8007>
- Wijgerde, T., et al. (2014). Coral calcification under daily oxygen saturation and pH dynamics reveals the important role of oxygen. *Biology Open*, 3(6), 489–493. <https://doi.org/10.1242/bio.20147922>